



SNF ratio, why and how?

June 21, 2022

Le Producteur de Lait Québécois Magazine

As of August 1, 2022, the maximum SNF ratio will drop from 2.30 to 2.25. This is an unsurprising development, given that the market needs a ratio of around 2.0. We could argue for a long time, going grocery shopping for consumers.



The demand for fat and protein is very strong. This is not the case for lactose. The lactose content of milk varies very little. To decrease the lactose production (in tons per year), you have to deliver a milk rich in fat first and then in protein. We will focus here on the fat content since it has the most influence in the SNF ratio calculation equation.

This article will not address in depth all the factors that influence the fat test of your herd. Ventilation, feeder space, water availability, feeding sequence, sorting, etc. are only a few of the many factors that can explain a fat test below expectations. We will limit ourselves to four technical parameters that are easy to measure and that can have important consequences on the results.

1. Genetics

The importance of genetics is undeniable. It is well understood today. The heritability for this trait is among the highest, around 50%. This means that it is an easy trait to improve compared to other traits such as conformation. In other words, a cow with a fat index of -0.50% mated to a

bull with +0.50% will produce a daughter with a genetic index close to the national average.

Knowing the genetic potential of the animals in your herd is of utmost importance. Strong genetics do not guarantee high butterfat in milk because there are many other factors involved. However, a herd with a genetic index of, say, -0.20% does not have what it takes to easily achieve a fat test of 4.0 kg/hl and better. This is not the case for the herd reported in Table 1. Its fat index (+0.11%) is well above the national average (+0.03%).

Table 1. Extract from the [Genetic Inventory Report](#) of a herd with a strong genetic background


GENETIC HERD INVENTORY				December 2021								
				HERD NUMBER	PAGE	TEST DATE						
				10 Dec 2021								
	Code	LPI	Pro\$	Prod Code	Milk kg	%RK	Fat kg	%RK	%	Protein kg	%RK	%
Cows	311	2901	1794		1018	75	52	79	0.11	46	85	0.10
Heifers (PA Prod)	280	3203	2596		1162	82	84	96	0.34	62	96	0.19
Total	591	3044	2174		1086		67		0.22	54		0.14
National Average		2508	579		318		15		0.03	12		0.02

Over the years, tools have been developed to help producers improve their control over the genetic development of their herd. Among these is the [Compass](#) application developed jointly by Holstein Canada and Lactanet.

Figures 1 and 2 show extracts for two herds. Which of the two herds do you think is in a better position to maintain an SNF ratio close to market needs? Herd A, of course. The deviation for fat of its recorded subjects

since 2016 (blue line) exceeds the 90-percentile rank for the breed. Herd B is now working on it. They are encouraged to continue.


Figure 1. Deviation for fat over time in herd A



Red line: national average

Figure 2. Deviation for fat over time in herd B

Fat Deviation



Red line: national average

2. Dry Matter Consumption in Forages

Since the cow is a ruminant, forage is the basis of her diet. A mixed ration (forages and concentrates) allows our cow to withstand higher production than just the need to feed her calf. We must go back to the basics. Forage is essential for rumination. Rumination is essential to the health of the microbial flora. The rumen microbial flora is essential to the cow's role on our planet, including the digestion of cellulose and hemicellulose to produce nutrients useful to humans. Most people are now familiar with [PROFILab](#).

[De novo fatty acids](#), measured in every milk sample in the tank, are the

very thing that allows you to judge the ruminal health of your cows.

We offer a simple way to evaluate the dry matter consumption in forages, i.e., kilograms per cow per day. That's all it takes. No complicated equations.

You will understand that for farms that handle chopped silage (vertical silos, horizontal silos, grinders, etc.), you will need the dry matter of the silage (moisture testers required).

For rations that include large bale silage (round or rectangular), we have a tool to calculate the weight of the bale in kilograms of dry matter. All you need is the bale size, the number of knives on the baler, the number of bales fed to lactating cows per week and the number of lactating cows.

Table 2. Examples of forage dry matter consumption calculation for a farm with chopped silage

Feed	% dry matter	kg per day	kg dry matter
Silage 1 st cut	40 %	15 kg	6.0 kg
Corn silage	35 %	20 kg	7.0 kg
Hay	90 %	1 kg	0.9 kg
Total		36 kg	13.9 kg

Producers who attended the Lactanet workshop on milk fat (*The Magic of a Milk Fat test*) last winter will remember that the target values (kg/day/cow) vary according to breed: 16 kg and more for Holsteins, 15 kg and more for Ayrshires, 18 kg for Brown Swiss, 13 kg and more for Jerseys and Canadian. The more corn silage we have in our ration, the higher the

target quantity.

If the dry matter consumption in forages is lower than the target values, the causes must be sought. The list is long. There is the [quality and conservation](#) of the forages, the management of the feeder, the sequence of the meals, the level of salt in the ration, etc. It is not always easy to find, but you should not stop looking until you have found the cause.

3. Effective Fibre

A herd with good genetics and exceptional forage dry matter intake may still be disappointed with the bulk tank fat test. For chopped grass silage, more than 10 factors influence the result and the theoretical chop length; forage adjustment is only one factor. The size of the swath plays a major role. In the field, chopping can be evaluated with the *Penn State* particle separator. The effective fibre is the particles found on the top two plates. In an article published in this magazine a few years ago, it was reported how the proportion of effective fibre varies from one silage to another. At Farm G (Figure 3), the rumination potential of the 2nd cut silage is much higher (73% vs. 50%) than that of the 1st cut.

Figure 3: Particle size assessment of 1st and 2nd cut silage from farm G

Ferme **Farm G**
Address

Analysis Date 2022-04-12

Sample Type **3**

1 = TMR, 2 = Corn Silage, 3 = Haylage

Data				
	Sample 1 Silage 1e		Sample 2 Silage 2e	
Screen	Weight (g)		Weight (g)	
Upper Sieve	22.0		65.0	
Middle Sieve	105.0		195.0	
Lower Sieve	116.0		85.0	
Bottom Plan	13.0		12.0	
Total	256.0		357.0	
Results				
Section 1. Particle Distribution				
	Sample 1 Silage 1e		Sample 2 Silage 2e	
Screen	Remaining Particles (% of total)	Cumulative (% under each sieve)	Remaining Particles (% of total)	Cumulative (% under each sieve)
Upper Sieve	9	91	18	82
Middle Sieve	41	50	55	27
Lower Sieve	45	5	24	3
Bottom Plan	5		3	
Section 2. Characterization of the sample				
	Sample 1		Sample 2	
Average Particle Length (mm)	6.69		9.91	
Standard Deviation (mm)	2.35		2.30	
Section 3. Recommended Particle Distribution				
	Sample Type: Haylage			
Screen	Remaining Particles (% of total)			
Upper Sieve	10 to 20			
Middle Sieve	45 to 75			
Lower Sieve	20 to 30			
Bottom Plan	Less than 5			



4. Milk/Concentrate Ratio

Another useful technical parameter to optimize your fat test is the milk/concentrate ratio. It is easy to calculate and gives a sounding board on several questions: well-balanced ration? Forage quality? Correct quantities of concentrates? Just to name a few.

In the Lactanet database, the milk to concentrate ratio has hovered around 3.30/1.0 since the winter of 2010-2011. The R farm profile (Table 3) is not the one we want. A ratio of 2.44 is too low. Now we have to investigate what could be the causes. In this case and at the time of the

visit to the farm, the dry matter consumption in forages was 12.3 kg and we remember that for Holsteins, we must aim for 16 kg.

Table 3. Calculation of the milk/concentrate ratio for farm R

Milk delivered (litres per day)	2,412 liters
Barley in the TMR (kg/day)	272 kg
Mineral in the TMR (kg/day)	12 kg
Robot feed (kg/day)	704 kg
Total concentrates (kg/day)	988 kg
Milk/concentrate ratio	2.44 liters/kg

There are many challenges in dairy production. From drought to input prices, the list goes on and on. The SNF ratio is another, but there are many tools to keep track of it. Don't forget that Lactanet is there to help you meet the needs of the market as best as possible, the key to the most profitable dairy production.

Action Plan for a Profitable Fat Test:

- Read the *Genetic Herd Inventory* report to know its genetic index for fat
- Use *Compass* to track the evolution of your genetic index for fat
- Calculate your forage dry matter consumption
- Use particle separators (e.g., *Penn State*) on a regular basis to determine the effective fibre of your forages - particles from the first two plates
- Calculate your milk/concentrate ratio

- Be aware of other factors that may influence the fat test – e.g., ventilation, feeder space, water availability, feeding sequence, sorting, etc.
- Do not hesitate to ask your technician or Lactanet advisor for help



By Jean Brisson, agr.

With his extensive knowledge in the field, Jean shares his expertise as a speaker and author of numerous publications on dairy production and contributes as a strategic advisor to the advancement of the dairy production industry.



By Diane Lequin agr.



By Valérie Martin agr.