



The milk fatty acid profile: What else can we learn?

October 14, 2020

Le Producteur de Lait Québécois Magazine

The fatty acid profile of bulk tank milk has been available to Quebec dairy producers since February 2020.

Each time milk is collected from the farm, farmers not only have access to regular milk composition (fat, protein, lactose and other solids) and herd somatic cell count, but can also learn more about the metabolism of their cows from urea and fatty acid analyses. The fatty acid profile (*de novo*, mixed and preformed) has helped many producers optimize herd management and diet, and we're now working to refine the use of these new indicators. Let's take a look at what we've learned so far, and what's on the horizon...

PROFILab in bulk tank milk: what we've learned

High variability between herds

American studies have reported that herds with higher *de novo* levels have shown better performances. Quebec data are, to some extent, consistent with that finding, showing a clear relationship between *de novo* fatty acids and bulk tank fat or protein (Figure 1). In fact, for every tenth of a unit increase in *de novo* fatty acids (0.10), fat content increases by two tenths (0.20). The link with true protein is also clear: for every 0.10 increase on the *de novo* scale, true protein increases by 0.12 units.

Figure 1: Each 0.1% increase in *de novo* fatty acids corresponds to a 0.2% increase in fat





Figure 1: 3387 herds in Quebec, 850 205 bulk tank results (April 2019-May 2020); Lactanet 2020

1

Our data for several thousand Quebec herds show, however, that the fatty acid profiles vary widely between farms. Breed is among the factors that have the greatest impact on the fatty acid. As shown in Figure 2, the levels of the different groups of fatty acids are higher in milk from predominantly Jersey herds, especially *de novo* and mixed fatty acids. More specifically, the *de novo* fatty acid content for Jersey herds is about 1.3 g/100 g of milk, as compared to around 1.05 g/100 g of milk for Holstein herds, with the other breeds falling in between. It should be noted that the variability between herds is high for all breeds, with the differences largely

attributable to ration composition, feeding behaviour and herd management.

Figure 2: Breed influences the fatty acid profiles of Quebec herds




High performance herds have average levels of *de novo* fatty acids

There is a common misconception that it is “normal” for high performance herds to have below-average *de novo* levels, due to a dilution effect related to above-average milk production. This is partly true, but the difference between high performance herds and average herds is minimal. To demonstrate this, Figure 3 shows the distribution of the fatty acid profiles of 3 300 conventional Holstein herds. The herds are categorized by annual production; the red box represents the herds in the bottom 10% for production, the orange box represents the herds that rank between the 10th and 80th percentile (26.6-37.9 kg of milk/cow/day), the green box represents the herds in the top 10% (37.9-40.7 kg of milk), and the blue box represents the top 2% of herds (over 40.7 kg of milk/cow/day). As the values below the graph indicate, the “blue” herds seem to have lower levels of *de novo* fatty acids in their milk, but the difference is about 0.01 g/100 g of milk on average, which is very close to the average presented in the PROFILab report.

Figure 3: High producing herds (kg/cow/day) make as many *de novo* fatty


acids as the average herd



Holstein herds only, December 2019 averages, categorized based on annual production; Lactanet 2020


If we repeat the exercise but categorize the herds according to their production in kg of fat/cow/day, the result is the same: the best herds have average *de novo* fatty acid contents (Figure 4). This graph shows that the variation between herds within a same production category is greater than the variation between the different categories. We are currently conducting a study of 100 Quebec herds to gain a better understanding of the factors behind this high variability, specifically those related to feeding, herd management and environment.

Figure 4: High fat producing herds (kg/cow/day) make as many *de novo* fatty acids as the average herd



How to interpret the boxplot?

A boxplot illustrates the variation within a group of data, in this case a group of herds. In the example below, the blue box represents 50% of herds, that is, all the herds that are “in the middle” of the group. The vertical lines extending below and above the box indicate the herds with the highest and lowest values.




PROFILab in bulk tank milk: what's next?

The bulk tank fatty acid profile has proven to be very useful for several producers to optimize feeding and herd management, or to get a better understanding of cow metabolism and performance. We will gain a more accurate understanding of this indicator as we use it more. It will undoubtedly become very interesting to use the fatty acid profile to predict a change in component levels, before the fat or protein levels actually change.

As an example, Figure 5 presents the profile of a herd in the summer of 2019, where we see that there were significant changes in the fatty acid profile in July and August. The first changes occurred in early July when body fat reserves were mobilized after a breakdown in the feeding equipment. The second change, on August 5, had a major impact on the fatty acid profile but didn't affect fat and protein levels. Later in August, a change of the same magnitude as in July, but longer lasting, led to an upheaval of the components and a drop in protein levels, which took several weeks to return to normal. Would this decline have been avoided with PROFILab? Perhaps, or it might have lessened the impact.

Figure 5



Lactanet 2020


5

Upcoming projects include a warning system indicating changes in the fatty acid profile that will enable producers and their service providers to react more quickly and prevent disastrous effects, and incorporating new and powerful analytical techniques such as machine learning. Also on the radar: predicting feed efficiency and methane production for lactating cows, and identifying biomarkers to certify grass-fed dairy herds. Stay tuned...

PROFILab individual cow: a glimpse at what's in the pipeline

At the same time, the rumen and the mammary gland are still adapting to lactation, and *de novo* fatty acids (synthesized in the mammary gland) are not being used to the full. Mixed fatty acids, a combination of the two previous sources, generally follow *de novo* fatty acids in early lactation. Based on the data that we've collected in the past few weeks in Quebec, and consistent with what's being seen in other parts of the world, most of the changes in the fatty acid profile seem to occur within the first 75 days of lactation, with the proportions stabilizing thereafter (Figure 6).

Figure 6: Milk fat composition is influenced by stage of lactation



Data from 199,388 Holstein cows in 3,106 herds; Lactanet 2020

It is clear that the changes in the different groups of fatty acids for cows in early lactation contain a great deal of valuable information, which can be used to ensure a successful transition and improve performance in our herds. Here are some specific examples:

1. Evaluate the mobilisation of body fat reserves

Cows in early lactation have a negative energy balance and very often use the energy stored in their body fat reserves to meet the extremely high demand for nutrients required to produce milk. Although this is a natural and well-known phenomenon, it is often difficult to estimate the extent of the mobilisation. The milk fatty acid profile can be used for that purpose. Moreover, in combination with the analysis of ketone bodies in milk (including β -hydroxybutyrate [BHB] measured by Ketolab or blood tests), the milk fatty acid profile

could provide greater insight into how cows manage this high demand for nutrients at the start of lactation, enabling producers to apply corrective measures to herd management and feeding during the transition period, as necessary.

2. Estimate weight loss in cows

As described in the section above, cows in early lactation mobilize body fat reserves, increasing the proportion of preformed fatty acids in the milk. By combining the composition of the different fatty acids and groups of fatty acids with other components such as total fat and BHB, it is possible to predict how much weight a cow will lose and respond appropriately to avoid excessive mobilisation that would diminish a cow's reproductive performance and potentially reduce longevity in the herd.

3. Assess the risk of ruminal acidosis

Measuring something that isn't visible to the naked eye, such as rumen pH for example, is definitely a way to use the milk fatty acid profile to benefit. This application is currently being validated through a joint project with researchers from Agrinova and Université Laval. If it proves possible to accurately predict rumen pH with the fatty acid analysis, it will then be possible to use the profile to assess the risk of acidosis, and possibly prevent more serious consequences by optimizing transition management, particularly for cows in early lactation.

Conclusion

The fatty acid profile has now been available for a number of months in Quebec, and new management tools and indicators are being developed and are expected to be launched in the coming months. Collectively, these tools based on the bulk tank or individual cow fatty acid analysis should enable us to refine our feeding and herd management practices to

maximize the efficiency and health of cows of all breeds, and ensure close follow-up for every cow in lactation in our herds.



By Débora Santschi, agr., Ph. D.

With her communication skills and passion for dairy production, Débora is a highly coveted speaker, both in Quebec and internationally. In addition to her role as Director of the R&D team at Lactanet, she is an associate professor in the departments of animal sciences at McGill University, Laval University, Manitoba and the Faculty of Veterinary Medicine (Université de Montréal).



By Daniel Warner